arXiv:astro-ph/0612366v1 14 Dec 2006

FULL TITLE
*ASP Conference Series, Vol. **VOLUME**, **YEAR OF PUBLICATION***
NAMES OF EDITORS

Balmer Absorption Lines in FeLoBALs

K. Aoki[1], I. Iwata[2], K. Ohta[3], N. Tamura[1], M. Ando[3], M. Akiyama[1],
G. Kiuchi[3], and K. Nakanishi[4]

Abstract. We discovered non-stellar Balmer absorption lines in two many-narrow-trough FeLoBALs (mntBALs) by the near-infrared spectroscopy with Subaru/CISCO. Presence of the non-stellar Balmer absorption lines is known to date only in the Seyfert galaxy NGC 4151, thus our discovery is the first cases for quasars. Since all known active galactic nuclei with Balmer absorption lines share characteristics, it is suggested that there is a population of BAL quasars which have unique structures at their nuclei or unique evolutionary phase.

1. Broad Absorption Line Quasar

Broad absorption line (BAL) quasars are quasars which show absorption lines in UV resonance lines, with velocity widths of $\sim 2000-20000$ km s^{-1}. BAL quasars are divided into three subtypes. High-ionization BAL quasars show absorption from C IV, N V, Si IV and Ly α. Low-ionization BAL quasars (LoBALs) show absorption from Mg II, Al III, and Al II, in addition to the above high-ionization absorption. A fraction of LoBALs shows absorption from Fe II and Fe III. They are called iron LoBALs (FeLoBALs).

Hall et al. (2002) discovered unique FeLoBALs which have tremendous number of absorption lines, so-called "many-narrow-trough FeLoBAL" (mntBAL). We have searched for mntBALs by visual inspection of $\sim$ 20000 spectra of quasars in $2.1 \leq z \leq 2.8$ and $1.3 \leq z \leq 1.65$ in the SDSS database. Hβ and Hα emission lines in these redshifts are expected to be redshifted into near-infrared regime. We have found 30 mntBALs on the redshift ranges.

2. Our results

We discovered Hα absorption line in SDSS 0839+3805 (Aoki et al. 2006) and Balmer-series absorption lines from Hα and H9 in SDSS 1723+5553 (Fig. 1). Presence of the non-stellar Balmer absorption lines is known to date only in the Seyfert galaxy NGC 4151, thus our discovery is the first cases for quasars. Recently Hall (2006) reports Balmer absorption lines in another FeLoBAL,

---

[1] Subaru Telescope, National Astronomical Observatory of Japan (NAOJ), 650 N. A'ohoku Place, Hilo, HI, 96720, U.S.A.

[2] Okayama Astrophysical Observatory, NAOJ, Okayama, 719-0232, Japan

[3] Department of Astronomy, Kyoto University, Kyoto, 606-8502, Japan

[4] Nobeyama Radio Observatory, NAOJ, Minamimaki Minamisaku, Nagano, 384-1305, Japan

SDSS J1259+1213. That quasar is also a possible mntBAL. The redshifts are determined to be 2.3179 and 2.1081 for SDSS 0839+3805 and SDSS 1723+5553, respectively, by Hα emission line. The Balmer absorption lines are blueshifted from the emission line by 520 and 5370 km s^{-1}, and the column density of neutral hydrogen is estimated to be 10^{18} and 5×10^{19} cm^{-2} for SDSS 0839+3805 and SDSS 1723+5553, respectively.

At least four active galactic nuclei (NGC 4151, SDSS 0839+3805, SDSS J1723+5553, and SDSS J1259+1213) have known to date show non-stellar Balmer absorption lines. Although they are much different in their luminosity, they share several characteristics. All four objects have Fe II absorption lines. NGC 4151 is not classified as FeLoBALs, however, its spectrum at the low luminosity phase shows many Fe II absorption lines (Kraemer et al. 2001). It is remarkably similar to SDSS 0839+3805 (Aoki et al. 2006). Strong [O III] emission line exists in all four objects. This attribute makes contrast to the previous observations of very weak or non-detection [O III] in LoBALs/FeLoBALs (Boroson & Meyers 1992; Yuan & Wills 2003). The FeLoBALs with Balmer absorption lines do not have strong Fe II optical emission lines. These facts suggest that the existence of such a population among BAL quasars, and they have unique structures at their nuclei or unique evolutionary phase.



Figure 1. Rest-optical spectrum of SDSS J1723+5553 in *JH*-band. Ordinate is a relative flux density in units of erg s^{-1} cm^{-2} μm^{-1} and abscissa is an observed wavelength in vacuum in micron. The rest wavelength is given along the top axis. The best fit is plotted as a thick solid line.

References

Aoki, K. et al. 2006, ApJ, 651, 84
Boroson, T. A., & Meyers, K. A. 1992, ApJ, 387, 442
Hall, P. B. et al. 2002, ApJS, 141, 267
Hall, P. B. 2006, astro-ph/0611922
Kraemer, S. B. et al. 2001, ApJ, 551, 671
Yuan, M. J., & Wills, B. J. 2003, ApJ, 593, L11